

Vaccination Recommendations for Broodmares- Previously Vaccinated Horses

Highly recommended or mandatory vaccines are underlined, suggested vaccines are not

Eastern and Western Encephalitis, Tetanus (EWT), West Nile Virus (WNV), Botulinum, Potomac recommended 4-6 weeks prior to foaling

EHV(Pneumabort/rhinoimmune)- 3,5,7, and 9 months out

Flu-vaccinate at least 2x yearly (depending on risk) with 1 dose 4-6 weeks before foaling

Rotavirus- 3 doses, at 8,9, and 10 months (each dose at least 4 weeks apart)

Strangles- Killed vaccine only -2x yearly with 1 dose 4-6 weeks before foaling

Rabies-yearly- some suggest 4-6 weeks prior to foaling to immunize the foal, others say the vaccine is not passed through the colostrums, and thus prior to breeding, or fall vaccination is recommended

****Do not vaccinate mares with more than 3 vaccines at any given time, and please leave 1 week between vaccination sets****

Example Vaccination Schedule:

3 months gestation: Pneumabort

5 months gestation: Pneumabort, Strangles, Flu

7 months gestation: Pneumabort

8 months gestation: Rotavirus, Rabies*

9 months gestation: Pneumabort, Rotavirus

6 weeks from due date: Strangles

5 weeks from due date: Flu, EWT, West Nile

4 weeks from due date: Potomac, Botulinum, Rotavirus